

New faunistic records of chironomids and phantom midges (Diptera, Chironomidae and Chaoboridae) from Ukraine indicate recent climatic refugia in the Eastern Carpathians

Peter Bitušík¹, Milan Novikmec², Marek Svitok^{2,3}, Ladislav Hamerlík^{1,4}

¹ Faculty of Natural Sciences, Matej Bel University, Tajovského 40, SK-974 01 Banská Bystrica, Slovakia

² Faculty of Ecology and Environmental Sciences, Technical University in Zvolen, T. G. Masaryka 24, SK-960 53 Zvolen, Slovakia

³ Department of Forest Ecology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Kamýcka 129, CZ-165 21 Prague, Czech Republic

⁴ Institute of Zoology, Slovak Academy of Sciences, Dúbravská cesta 9, SK-845 06 Bratislava, Slovakia

Corresponding author: Peter Bitušík (peter.bitušik@umb.sk)



Academic editor: Viktor Baranov

Received: 15 April 2024

Accepted: 28 June 2024

Published: 9 September 2024

ZooBank: <https://zoobank.org/E6BB88C5-5795-4E80-9888-82146DD4CE9C>

Citation: Bitušík P, Novikmec M, Svitok M, Hamerlík L (2024) New faunistic records of chironomids and phantom midges (Diptera, Chironomidae and Chaoboridae) from Ukraine indicate recent climatic refugia in the Eastern Carpathians. ZooKeys 1211: 349–367. <https://doi.org/10.3897/zookeys.1211.125436>

Copyright: © Peter Bitušík et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Abstract

The aquatic insect fauna of the Eastern Carpathians is poorly known, especially in Ukraine. To address this knowledge gap, we conducted faunistic surveys of Chironomidae and Chaoboridae in 2018 and 2021. The study involved sampling of 11 watercourses and 10 mountain lakes situated in the Ukrainian part of the Eastern Carpathians. A total of 101 taxa were identified, including 40 chironomid species and one genus that have been recorded for the first time from Ukraine. The occurrence of one species previously considered as “doubtfully present” in Ukraine was confirmed by this study. One of the two identified phantom midge species, *Chaoborus* (s. str.) *obscuripes* (van der Wulp, 1859), is recorded for the first time from Ukraine. The most intriguing records are chironomid species *Cricotopus* (s. str.) *beckeri* Hirvenoja, 1973, *Eukiefferiella bedmari* Vilchez-Quero & Laville, 1987, and *Pseudorthocladius* (s. str.) *berthelemyi* Moubayed, 1990. These species have Mediterranean distribution and their occurrence in the Eastern Carpathians could be remains of once-widespread populations that currently survive in the Carpathian refugia due to adverse climatic conditions in the former distribution area. The high number of first records from a relatively small number of sites indicates a great gap in the knowledge of the Ukrainian chironomid fauna.

Key words: climatic relicts, mountain lakes, pupal exuvia, submontane rivers

Introduction

Chironomids are the most ubiquitous free-living holometabolous insects known from all zoogeographic regions, and all climatic zones from the tropics to the polar regions, including Antarctica (Ashe and O'Connor 2009). Recently, 7290 species belonging to nearly 440 genera and 11 subfamilies have been described worldwide (Ferrington 2007; Pape et al. 2011). In immature stages, most species inhabit various types of freshwaters although some species thrive in brackish water and intertidal pools, and few are truly marine. Finally, semi-terrestrial and fully terrestrial species are also known (Sæther et al. 2000). Among aquatic insects, Chironomidae is the most species-rich insect family

found in freshwater ecosystems (Cranston 1995; Ferrington 2007). The species richness of the family recorded from only one stream locality is often astonishing. In some cases, the number of chironomid species recorded is higher than the diversity of all the other benthic macroinvertebrates (own observations).

Chironomidae can withstand an extremely wide range of environmental conditions in terms of water column depth, temperature, pH, dissolved oxygen, habitat drying and, finally, the gradient of human impacts such as pollution, habitat modification, and changes in watersheds (Ferrington 2007 and references therein). Consequently, chironomids have attracted the attention of researchers around the world as biological indicators for environmental impact assessments, ecosystem health, palaeolimnological reconstructions, and climate change (Resh and Rosenberg 2008; Eggermont and Heiri 2012; Nicacio and Juen 2015 and references therein). Compared to chironomids, the global diversity of phantom midges is low. The Chaoboridae family consists of about 51 extant species in six genera and two subfamilies (Borkent 2014). The immature stages usually live in standing waters, in some cases in small, temporary ponds. Larvae are predators and mostly planktonic; they are often considered keystone species that can eliminate or strongly suppress other invertebrates in the community (MacKay et al. 1990). Subfossil remains of the *Chaoborus* genus have been used in palaeoenvironmental research but also in contemporary ecological studies (e.g., Luoto and Nevalainen 2009; Tolonen et al. 2012). Despite the indisputable importance of both the chaoborids and chironomids and the rapid progress in the knowledge of their distribution, there are still areas that are “terra incognita”. The Eastern Carpathians are undoubtedly one of them.

Here, we present results from an ongoing faunistic inventory of chironomids and chaoborids from Ukrainian Carpathian Mountain lakes, which are supplemented by results from earlier investigations of the flowing waters in this territory.

Materials and methods

Study area

This study was conducted in the Ukrainian Carpathians located in the northern part of the Eastern Carpathians, extending through the western part of Ukraine (Fig. 1). The total length of this mountain range is approximately 240 km with total area of ca 24,000 km². The Ukrainian Carpathians are medium-high mountains with the highest elevation slightly exceeding 2000 m a.s.l. (Novikoff and Hurdu 2015; Vyshnevskyi and Donich 2021). The studied area is characterised by complex geology consisting mostly of flysch with different constituents. Only small areas are formed by limestone, shales, and volcanic rocks, predominantly andesites and gneisses (Ivanik et al. 2019). We sampled 11 streams and 10 mountain lakes (Table 1). Except for lakes Sinevir and Dragobratske located in deciduous and coniferous forests, respectively, all studied lakes are situated above 1500 m a.s.l., i.e., above formerly accepted climatic tree line (Kobiv 2017). The semi-natural meadows and pastures (so-called *polonynas* in the vernacular) dominate the catchment vegetation. The proportion of dwarf pine (*Pinus mugo*), dwarf juniper (*Juniperus communis* subsp. *nana*), green alder (*Alnus viridis*), and scattered Norway spruce (*Picea abies*) is different in individual catchments.




Figure 1. Map showing of sampling localities of Chironomidae and Chaoboridae in the Ukrainian part of the Eastern Carpathians. The site abbreviations correspond to the Table 1. Inset shows location of the investigated territory within the Carpathian Mountains (grey area).

Table 1. Basic characteristics of the sampling sites. Strahler stream order was estimated from Google Earth Pro. Stream width, depth and lake depth were estimated in the field. Lake area was derived from Google Earth Pro using Polygon tool.

Stream/Lake name	Code	Latitude, Longitude	Elevation (m)	Stream order	Av. width (m)	Av. depth (m)
Zhdenivka River	S1	48°46.29'N, 22°58.68'E	409	IV.	6	0.30
Nameless headwater stream	S2	48°49.19'N, 22°58.58'E	777	II.	2	0.15
Tributary of the Zhdenivka River	S3	48°47.98'N, 22°57.41'E	454	III.	3	0.20
Latoricja River	S4	48°45.00'N, 23°00.75'E	360	IV.	20	0.30
Gluhana peat bog channel	S5	48°27.94'N, 23°37.94'E	596	I.	0.4	0.10
Rika River (village Soimy)	S6	48°34.00'N, 23°28.53'E	449	IV.	35	0.15
Rypenka River (village Soimy)	S7	48°33.92'N, 23°28.29'E	448	IV.	25	0.30
Rika River (above Zaperedillia village)	S8	48°29.08'N, 23°30.24'E	382	V.	30	0.50
Nameless tributary of the Volovets River	S9	48°33.14'N, 23°35.51'E	725	II.	2	0.15
Tereblia River (village Sinevyr)	S10	48°28.25'N, 23°38.16'E	573	IV.	18	0.30
Sukhar brook (village Kolochava)	S11	48°25.40'N, 23°42.58'E	580	III.	10	0.30
Tisa River (Vinohradiv)	S12	48°08.06'N, 23°05.15'E	124	VII.	120	–
				Area (ha)	Max. depth (m)	
Sinevir Lake	L1	48°37.01'N, 23°41.04'E	989	4.20	22	
Brespo Lake	L2	48°08.62'N, 24°30.94'E	1627	0.03	0.4	
Bolotnoe ozerce	L3	48°08.55'N, 24°31.25'E	1695	0.01	0.9	
Brebeneskul Lake	L4	48°06.10'N, 24°33.74'E	1793	0.60	3.2	
Dragobratske Lake	L5	48°14.45'N, 24°14.45'E	1382	0.07	1.5	
Apshynets Lake	L6	48°16.91'N, 24°09.53'E	1491	2.97	1.8	
Geryshaska (Dohiaska) Lake	L7	48°16.25'N, 24°09.93'E	1585	2.58	2	
Kosivske Lake	L8	48°15.78'N, 24°11.96'E	1614	0.13	1	
Ivor Lake	L9	48°13.70'N, 24°14.10'E	1606	0.04	2	
Small Ivor Lake	L10	48°13.69'N, 24°14.07'E	1602	0.04	0.6	

All the studied flowing water sites except the Tisa River were located in the Carpathians. Most of the sampling sites were situated in valleys of the sub-montane belt up to 550–600 m a.s.l. (established according to climate and vegetation characteristics; Golubets 1977), only two sites (S2, S9) were situated in the lower montane belt above 700 m a.s.l. Stream bottom substrates were mostly dominated by cobbles with an admixture of finer components such as gravel and sand. The bottom habitats of both mountain brooks were more complex due to the presence of boulders and a considerable proportion of woody debris. Because of the bed surface roughness and shallow depth, turbulent flow is generally prevailing. Even though most sites were located near roads and often in settlements, the channels, banks, and riparian vegetation were visibly well preserved, without the signs of artificial modifications.

The above characteristics mostly do not apply to the Upper Tisa River. The river stretch close to the town of Vinohradiv flows in a lowland landscape (the Hungarian lowland ecoregion; Afanasyev et al. 2020). The floodplain is covered with alluvial forests containing willow and poplar. In the studied section, the low slope and high discharge altered the river channel morphology significantly, forming a high degree of channel sinuosity. The basic characteristics of the sampling sites are summarised in Table 1.

Sampling and identification

Chironomids and chaoborids were collected during the sampling campaigns in May 2018 (flowing waters and Sinevir Lake) and in August 2021 (mountain lakes in the Chornohora and Svidovets Massifs). A hand net attached to a telescopic handle (mesh size 250 µm, frame diameter 25 cm) was used to skim the water surface and collect floating material along the shores of streams and lakes. In lakes, the material was collected at the leeward shore; in flowing waters, the floating material was collected along an ~ 100-meter-long stretch while moving upstream.

On the shore, the netted sample was placed in a labelled 100-ml plastic bottle and preserved with 75% ethanol. In the laboratory, the samples were placed in a Petri dish and all chironomid material was picked up under a stereomicroscope (7.5–50×). Sorted exuvia, pupae, and adults were mounted on microscopic slides and identified following the keys of Langton and Visser (2003), Ekrem (2004), Stur and Ekrem (2006), and Langton et al. (2013) for pupal exuvia and Langton and Pinder (2007a, 2007b) for adults under high magnification (400×) using phase contrast. Chaoborid larvae, which were collected accidentally during the chironomid sampling of lakes, were identified using Parma (1969) and Sæther (1997).

The nomenclature and distribution of species are consistent with Fauna Europaea (de Jong 2016), and Ashe and O'Connor (2009, 2012). Voucher specimens are deposited in the collections of the Dept. of Biology and Environmental Studies, Faculty of Natural Sciences, Matej Bel University in Banská Bystrica.

Results and discussion

A total of 2088 specimens were collected and identified as 99 chironomid species/taxa (belonging to 43 genera from 5 subfamilies) and two chaoborid species of the same subfamily and genus. Altogether, 40 species and one genus

of Chironomidae, and one species of Chaoboridae were recorded for the first time in Ukraine. The occurrence of one chironomid species, *Nilotanypus dubius* (Meigen, 1804), previously considered as “doubtfully present” in Ukraine (Spies and Sæther 2013) was finally confirmed. A list of all species/ taxa recorded is given below (sampling site codes refer to Table 1; **Pe** after the genus name refers to a morphotype not associated with an adult by Langton (1991); # – previously doubtfully present in Ukraine, * – first record of species/ genus from Ukraine). For detailed information on the abundance and life stages of collected specimens see Supplementary file 1.

CHIRONOMIDAE

Tanypodinae

Procladius (*Holotanypus*) *choreus* (Meigen, 1804): L5, L6, L9

Nilotanypus dubius (Meigen, 1804)#: S1, S6, S7, S8

Thienemannimyia carnea (Fabricius, 1805)*: S7

Zavrelimyia barbatipes (Kieffer, 1911): L6

Diamesinae

Diamesa (*Diamesa*) *cinerella* Meigen, 1835*: S8

Diamesa (*Diamesa*) cf. *tonsa* (Haliday, 1856): S6

Diamesa (*Diamesa*) *vaillanti* Serra-Tosio, 1972*: S7

Potthastia Pe1 Langton, 1991: S4, S7, S8

Sympotthastia macrocera Serra-Tosio, 1973*: S6

Prodiamesinae

Prodiamesa olivacea (Meigen, 1818): L1

Orthoclaadiinae

Brillia bifida (Kieffer, 1909): S3

Brillia flavifrons (Johannsen, 1905)*: S7

Corynoneura celtica Edwards, 1924*: S1, S3, S10

Corynoneura cf. *scutellata* Winnertz, 1846: L1, L6

Corynoneura Pe2a Langton, 1991: S1, S4, S5, S6, S7, S8

Corynoneura Pe4 Langton, 1991: S1

Cricotopus (*Cricotopus*) *annulator* Goetghebuer, 1927: S1, S4, S5, S6, S7

Cricotopus (*Cricotopus*) *beckeri* Hirvenoja, 1973*: S4, S6, S8

Cricotopus (*Cricotopus*) *curtus* Hirvenoja, 1973*: S6

Cricotopus (*Cricotopus*) *fuscus* (Kieffer, 1909): S1, S11

Cricotopus (*Cricotopus*) *pallidipes* Edwards, 1929*: S8

Cricotopus (*Cricotopus*) *similis* Goetghebuer, 1921*: S1, S7, S8

Cricotopus (*Cricotopus*) *tremulus* (Linnaeus, 1758)*: S1, S3, S6, S8

Cricotopus (*Cricotopus*) *trifascia* Edwards, 1929: S4, S11

Cricotopus (*Cricotopus*) *vierriensis* Goetghebuer, 1935: S1, S4, S6, S7, S8

Cricotopus Pe 17 Langton, 1991: S4, S8

Cricotopus (Isocladius) reversus Hirvenoja, 1973*: L1
Cricotopus (Isocladius) sylvestris (Fabricius, 1794): L6, L7
Cricotopus (Isocladius) Pe 5 Langton, 1991: L6, L7
Cricotopus (Paratrichocladius) rufiventris (Meigen, 1830): S4, S7, S10
Eukiefferiella bedmari Vilchez-Quero & Laville, 1987*: S4
Eukiefferiella brevicealcar (Kieffer, 1911): S2, S3
Eukiefferiella clypeata (Thienemann, 1919)*: S6, S7, S11
Eukiefferiella coerulescens (Kieffer, 1926): S3, S8
Eukiefferiella devonica (Edwards, 1929): S9
Eukiefferiella fuldensis Lehmann, 1972: S10
Eukiefferiella ilkleyensis (Edwards, 1929): S4, S6, S7, S8, S10
Euryhapsis Pe1 Langton, 1991: S11
Heleniella serratosioi Ringe, 1976*: S1, S4, S6, S7, S8, S9, S10, S11
Krenosmittia boreoalpina (Goetghebuer, 1944)*: S1, S3, S6, S7, S8, S10
Krenosmittia camptophleps (Edwards, 1929)*: S10
Limnophyes cf. asquamatus Andersen, 1937: L3
Nanocladius (Nanocladius) parvulus (Kieffer, 1909): S3, S4, S6, S8
Nanocladius (Nanocladius) rectinervis (Kieffer, 1911)*: S6, S7, S8, S10, S11
Orthocladius (Orthocladius) dentifer Brundin, 1947: L7
Orthocladius (Orthocladius) excavatus Brundin, 1947*: S7, S8, S11
Orthocladius (Orthocladius) oblidens (Walker, 1856)*: S11
Orthocladius (Orthocladius) pedestris Kieffer, 1909*: S1, S3, S4, S7, S8, S11
Orthocladius (Orthocladius) rivinus Potthast, 1914*: S3
Orthocladius (Orthocladius) rubicundus (Meigen, 1818): S1, S3, S4, S7, S8, S10
Orthocladius (Euorthocladius) ashei Soptonis, 1990*: S4, S7, S8, S11
Orthocladius (Euorthocladius) rivicola Kieffer, 1911: S3, S6, S8, S10, S11
Orthocladius (Euorthocladius) rivulorum Kieffer, 1909: S4, S6, S7, S8
Paracricotopus niger (Kieffer, 1913)*: S4, S6, S7, S8, S11
Parakiefferiella bathophila (Kieffer, 1912)*: S4, S7, S8
Parametriocnemus stylatus (Spaerck, 1923): S1, S3, S4, S6, S7, S8, S9, S10, S11
Psectrocladius (Psectrocladius) limbatellus (Holmgren, 1869): S11
Psectrocladius (Psectrocladius) oligosetus Wuelker, 1956: L2, L3, L9, L10
Psectrocladius (Psectrocladius) schlienzi Wuelker, 1956*: L1
Pseudorthocladius (Pseudorthocladius) berthelemyi Moubayed, 1990*: S6
Rheocricotopus (Psilocricotopus) chalybeatus (Edwards, 1929): S1, S4, S6, S7, S8
Rheocricotopus (Rheocricotopus) fuscipes (Kieffer, 1909): S1, S3, S7, S8
Rheosmittia spinicornis (Brundin, 1956)*: S2, S3, S10
Symbiocladius rhithrogenae (Zavrel, 1924): S11
Synorthocladius semivirens (Kieffer, 1909): S3, S10
Thienemanniella majuscula (Edwards, 1924): S1
Thienemanniella Pe 1b Langton, 1991: S3
Thienemanniella Pe 2 Langton, 1991: S10
Tvetenia verralli (Edwards, 1929)*: S6

Chironominae

Benthalia sp.: L7
Demicryptochironomus Pe1 Langton, 1991: S11
Microtendipes chloris (Meigen, 1818): S5, S7

Microtendipes pedellus (De Geer, 1776): L6
Paracladopelma mikianum (Goetghebuer, 1937)*: S6, S11
Phaenopsectra flavipes (Meigen, 1818): S5, L1, L2, L6, L7
Polypedilum (*Polypedilum*) *albicorne* (Meigen, 1838)*: S4, S6, S7, S11
Polypedilum (*Polypedilum*) *laetum* (Meigen, 1818): S11
Polypedilum (*Polypedilum*) *nubeculosum* (Meigen, 1804): S11
Polypedilum (*Pentapedilum*) *sordens* (van der Wulp, 1875): S7
Polypedilum (*Pentapedilum*) cf. *uncinatum* (Goetghebuer, 1921): L2, L5, L7
Polypedilum (*Tripodura*) cf. *apfelbecki* (Strobl, 1900): S6
Polypedilum (*Uresipedilum*) *convictum* (Walker, 1856): S4, S11
Cladotanytarsus (*Cladotanytarsus*) *atridorsum* Kieffer, 1924: L6, L7, L8
Cladotanytarsus (*Cladotanytarsus*) *vanderwulpi* (Edwards, 1929): S6, S7, S8
Micropsectra atrofasciata (Kieffer, 1911): S6, S7
Micropsectra lindrothi Goetghebuer, 1931*: L7
Micropsectra lindebergi Saewedal, 1976/ *insignilobus* Kieffer, 1924: S5
Neozavrelia Pe1 Langton, 1991*: S1, S4, S5, S6, S7, S8
Paratanytarsus austriacus (Kieffer, 1924): L6
Paratanytarsus dissimilis (Johannsen, 1905)*: S5
Paratanytarsus laccophilus (Edwards, 1929): L2, L5, L7, L8, L9, L10
Rheotanytarsus pentapoda (Kieffer, 1909)*: S6, S7
Rheotanytarsus rhenanus Klink, 1983*: S1, S6, S7
Stempellinella flavidula (Edwards, 1929)*: S8
Tanytarsus aberrans Lindeberg, 1970*: L6
Tanytarsus debilis (Meigen, 1830)*: L5, L7
Tanytarsus gregarius Kieffer, 1909: L2, L4
Tanytarsus heusdensis Goetghebuer, 1923*: S6, S7
Virgatanytarsus Pe1 Langton, 1991: S4

CHAOBORIDAE

Chaoborinae

Chaoborus (*Chaoborus*) *crystallinus* (De Geer, 1776): L3
Chaoborus (*Chaoborus*) *obscuripes* (van der Wulp, 1859)*: L7

Of the 99 recorded chironomid taxa, 22 were found exclusively in lakes, and 20 of them only in alpine lakes. Fourteen lacustrine species/ taxa (i.e., 70%) were also found in lakes during our previous research (Bitušík et al. 2020) indicating a similar species composition of all the studied lakes. Phantom midges were not deliberately targeted since the larvae recorded were caught by chance while collecting pupal exuvia. However, one of the species recorded, *Chaoborus* (s. str.) *obscuripes* (van der Wulp, 1859), appears to be the first record in the Ukrainian Carpathian alpine lakes. Most chironomid species have been found in flowing waters. Overall, they are considered typical for streams and rivers of Western and Central Europe (e.g., Caspers 1991; Laville and Vinçon 1991; Ruse 1995; Orendt 2002a, 2002b; Schöll and Haybach 2004; Bitušík et al. 2006; Calle-Martínez and Casas 2006; Prat et al. 2016 and citation therein). Because most of the newly recorded species are widespread in stagnant and flowing waters in Europe and the Palaearctic Region, and do not increase our

knowledge on their ecology, only species with restricted distributions and rarely collected elsewhere are discussed further in more detail. Regarding the common species not discussed further in the text, their geographical distribution is documented in Ashe and O'Connor (2009, 2012) and de Jong (2016), while their ecology is summarised in Moller Pillot (2009, 2013) and Vallenduuk and Moller Pillot (2007).

Family Chironomidae
Subfamily Diamesinae
Tribe Diamesini

***Diamesa (Diamesa) vaillanti* Serra-Tosio, 1972**

Material examined. 1 pupal exuvium, Rypenka river (S7), 7 May 2018.

Distribution. Palaearctic: Germany, Switzerland, France, Italy, Austria, Slovakia, Poland, Russia, Spain, Turkey, Morocco (Ashe and O'Connor 2009), and Azerbaijan (Kownacki 1985).

Habitat. Rheophilic species inhabiting high-altitude springs, streams (including glacier-fed), and rivers with rocky bottoms, but also alpine lakes (Bitušík 2004; Lencioni et al. 2011; Rossaro and Lencioni 2015). The species is cold-stenothermal (e.g., Rossaro 1991) although the findings from streams in the High Atlas indicate that it can tolerate relatively high temperatures of up to 22 °C (Azzouzi et al. 1992).

Remarks. The occurrence of the species is limited to waters at high altitudes. Since our record is from 448 m a.s.l., our finding is exceptional.

***Sympotthastia macrocera* Serra-Tosio, 1973**

Material examined. 2 pupal exuvia, Rika River (S6) in Soimy village, 7 May 2018.

Distribution. Palaearctic. For a long time, known only from Western Europe (France, Germany; Ashe and O'Connor 2009). More recent data comes from the Drava River in Croatia (Kresonja 2018). Our record is evidence for the current easternmost occurrence in Europe, but there are indications that the distribution of this species may extend as far as the Ural Mountains (Krashenninnikov 2012).

Habitat. Generally, larvae of the genus *Sympotthastia* inhabit cold running waters and springs (Sæther and Andersen 2013). A few data indicate that *S. macrocera* is probably a rheophilic species.

Remarks. *Sympotthastia macrocera* appears to be a relatively rare species with little known ecology.

Subfamily Orthocladiinae

***Cricotopus (Cricotopus) beckeri* Hirvenoja, 1973**

Material examined. 1 pupal exuvium, Latoricja River (S4), 5 May 2018; 1 pupal exuvium, Rika River (S6); 2 pupal exuvia, Rika River (S8) above Zaperedillia village, 7 May 2018.

Distribution. Palaearctic. France, Spain, Greece, Madeira, Corsica, Turkey, Algeria, Morocco, Slovakia. Its questionable occurrence in Finland (Ashe and O'Connor 2012) was not accepted later Paasivirta (2014).

Habitat. Principally inhabits the rhithral zone of streams at lower altitudes (Hirvenoja and Moubayed 1989; Kettani and Langton 2012; Moubayed-Breil and Ashe 2016).

Remarks. *Cricotopus beckeri* has been considered an exclusively Mediterranean species (Laville and Reiss 1992). Our record is the second reliable finding in the Carpathians (Bitušík and Langton 1994) far from its continuous distribution. We assume that the isolated populations in the Carpathians could be remnants of once widespread populations, which currently survive in refugia due to adverse climatic conditions. Thus, they could be considered a climatic relict, as customary for some plant species (Molnár et al. 2017). Interestingly, Reiss (1986) already hypothesised that the extra-Mediterranean occurrence of another Mediterranean chironomid, *Paratanytarsus mediterraneus* Reiss, 1981, could have a relict character in the Middle Rhine.

***Cricotopus (Cricotopus) pallidipes* Edwards, 1929**

Material examined. 1 pupal exuvium, Rika River (S8), 7 May 2018.

Distribution. Palaearctic. Finland, Norway, France, Portugal, Spain, Germany, Great Britain, Ireland, Romania, Hungary, Russia, Lebanon, and Morocco (Cobo et al. 2002; Soriano and Cobo 2006; Móra et al. 2006; Ashe and O'Connor 2012).

Habitat. The ecological requirements of this species are still unclear. It has been found in flowing and stagnant waters in cold climatic zones (Aagaard et al. 1997; Pozdeev 2012) to warm rivers, canals, lakes, and marshes in central and southern Europe, and North Africa (e.g., Laville and Tourenq 1968; Móra et al. 2006; Abbou and Fahde 2017; Moubayed et al. 2019). French authors (Tourenq 1976; Moubayed et al. 2019) consider it a lacustrine species, tolerant of low oxygen content.

Remarks. Currently known only from few European countries. It does not seem to be abundant anywhere. In Bavaria and the Sauerland Mountains (Germany), it is listed among possibly endangered species; however, its status is unknown (Orendt and Reiff 2003; Dittmar 2012).

***Eukiefferiella bedmari* Vilchez-Quero & Laville, 1987**

Material examined. 3 pupal exuvia, Latoricja River (S4), 5 May 2018.

Distribution. Palaearctic. France, Spain, Greece, Corsica, Turkey, Lebanon, Algeria, and Morocco (Ashe and O'Connor 2012).

Habitat. Streams and rivers (Laville and Langton 2002; Chaib et al. 2013; Moubayed-Breil and Ashe 2016).

Remarks. *E. bedmari* is a circum-mediterranean faunistic element (Laville and Reiss 1992; Moubayed-Breil 2008). Our unexpected extra-Mediterranean finding suggests the relict character of its population in the Carpathians (see comments to *Cricotopus beckeri*).

***Orthocladius (Orthocladius) rivinus* Potthast, 1914**

Material examined. 1 pupal exuvium, left-hand tributary of Zhdenivka River (S3), 7 May 2018.

Distribution. Palaearctic. Norway, Great Britain, Ireland, Austria, Slovakia, Hungary, Poland, Belarus, Germany, Switzerland, France, Italy, Spain, Canary Islands, and Portugal (Cobo et al. 2002; Ashe and O'Connor 2012; Moller Pillot 2013; Móra et al. 2013; Sołtys-Lelek et al. 2014).

Habitat. Rheophilic species inhabiting springs and flowing waters from small streams to large rivers, although it has been reported also from lakes (Langton and Visser 2003). Rossaro et al. (2003) underline its preference for cold waters, but some findings question this (e.g., Langton and Orendt 1996; Móra et al. 2013).

Remarks. The species is known from a few European countries and is generally considered rare. Like the ambiguous data on its ecology, this may also be the result of misidentification.

***Psectrocladius (Psectrocladius) schliezi* Wuelker, 1956**

Material examined. 1 pupal exuvium, Lake Sinevir (L1), 7 May 2018.

Distribution. Palaearctic? In addition to some European countries (Austria, Denmark, Finland, Germany, Great Britain, Italy, Moldova, Netherlands, Norway, Portugal, Slovakia, Czech Republic, Spain, Sweden, Switzerland; Ashe and O'Connor 2012; Syrovátka and Langton 2015), it was also recorded in Mongolia (Hayford 2005). However, the species may have a Holarctic distribution, provided that its presence in North America can be confirmed (Sealock and Ferrington 2008). Baranov et al. (2024) found a species resembling *P. schliezi* in Namibia, but it is possible that the specimen belongs to a yet undescribed species of *Psectrocladius* related to *P. schliezi*.

Habitat. Different types of stagnant waters from lakes to pools. For example, the only record from the Carpathians comes from a shallow pond in an exploited part of an alkaline fen (Bitušík and Illéšová 1998). Its occurrence in slowly flowing waters is exceptional (de Beauvesère-Storm and Tempelman 2009).

Remarks. The records are scattered across Europe, and it seems that the species is not abundant anywhere (Moller Pillot 2013).

***Pseudorthocladius (Pseudorthocladius) berthelemyi* Moubayed, 1990**

Material examined. 11 pupal exuvia, Rika River (S6), 7 May 2018.

Distribution. Palaearctic. Austria, Bulgaria, Corsica, France, Germany, Portugal, Slovakia, Spain, Turkey, and Morocco (Ashe and O'Connor 2012).

Habitat. Mountain streams and rivers with stony bottoms. The species is rheophilic, cold-stenothermal with high demand for dissolved oxygen (Martínez et al 1995; Moubayed-Breil et al. 2012). It can also inhabit hygropetric sites (Moubayed-Breil 2008).

Remarks. The species is considered a Mediterranean element (Moubayed-Breil 2008) with an originally circum-mediterranean distribution

(Laville and Langton 2002). The extra-Mediterranean occurrence in more northerly countries indicates its relict character (see comments to *Cricotopus beckeri*).

Subfamily Chironominae

Tribe Chironomini

Paracladopelma mikianum (Goetghebuer, 1937)

Material examined. 2 pupal exuvia, Rika River (S6), 7 May 2018; 1 pupal exuvium, Tisa River (S12), 8 May 2018.

Distribution. Palaearctic. The species was recorded only from a few countries in Europe (e.g., Spain, Hungary, Slovakia, Portugal, France, Germany, Romania) and North Africa (Morocco, Lebanon).

Habitat. It is a rheophilic species inhabiting fast-flowing streams and rivers (Moller Pillot 2009). Although Calle-Martínez and Casas (2006) listed the species in a chironomid community usually associated with low-temperature or torrential mountain streams, our finding from Tisa River and other records from large lowland rivers (Gandouin et al. 2006; Klink 2010) indicate that the species does not have as strict cold temperature preferences as though previously (Ringe 1974).

Remarks. Laville and Vinçon (1986) considered the species to be a Mediterranean-Palaearctic element whose northern limit is located in the Pyrenees, the Alps, and the Carpathians.

Tribe Tanytarsini

Neozavrelia Goetghebuer, 1941

Material examined. 11 pupal exuvia, Zhdenivka River (S1), 5 May 2018; 17 pupal exuvia, Latoricja River (S4), 5 May 2018; 1 pupal exuvium, Rika River (S6), 7 May 2018; 18 pupal exuvia Rika River (S8), 7 May 2018; 19 pupal exuvium, Rypenka River (S7), 7 May 2018; 1 pupal exuvium, channel at Gluhana peat bog (S5), 7 May 2018.

Distribution. Species-rich genus (38 valid species, de Jong 2016) with a worldwide distribution except for Africa and Neotropics (Epler et al. 2013). Five species have been recorded in Europe, three of which are reliably confirmed in the Carpathians: *N. improvisa* Fittkau, 1954, *N. luteola* Goetghebuer, 1941 (Giłka 2007), and *N. cuneipennis* (Edwards, 1929) (Tatole 2023).

Habitat. Larvae of *Neozavrelia* inhabit streams, rivers, lakes, and ponds in peat bogs; they are also known from hygropetric sites, and one species lives in a hot spring (Epler et al. 2013).

Remarks. Except for *N. cuneipennis* (= *N. longappendiculata* Albu, 1980), the morphological characteristics of the pupae do not yet allow for distinguishing the European species (Langton and Visser 2003). The morphotype *Neozavrelia* Pe1 Langton, 1991 includes four species: *N. bernensis* Reiss, 1968, *N. fuldensis* Fittkau, 1954, *N. improvisa*, and *N. luteola*.

Family Chaoboridae

Chaoborus (Chaoborus) obscuripes (van der Wulp, 1859)

Material examined. 1 larva, Lake Geryshaska (L7), 15 September 2021.

Distribution. Palaearctic. The species is widespread mainly in Northern and Western Europe, but also in Poland and the European part of Russia (Borkent 1981; de Jong 2016).

Habitat. Small, shallow nutrient-poor, meso- and polyhumic ponds with pH 4.5–5.5 (Nilssen 1974; Joniak and Domek 2006; Kuper and Verberk 2011), often fishless. Larger larvae with darker pigmentation are more sensitive to visually dependent predators (Stenson 1981).

Remarks. The species seems to occur sporadically and mostly in small numbers (Borkent 1981), which is probably related to its ecological requirements for water chemistry and the absence of fish.

The first annotated checklist of Ukrainian Chironomidae consists of 302 species (Baranov 2011a). However, this list requires revision because it contains invalid species identified solely on the basis of larvae using outdated identification keys. In recent decades, the study of taxonomy, ecology, and biogeography of chironomids in Ukraine has intensified (Baranov 2011b, 2013, 2014; Baranov and Przhiboro 2014; Baranov and Ferrington 2013; Moubayed-Breil and Baranov 2018; Didenko et al. 2021). Our survey revealed a significant gap in the taxonomic knowledge of Ukrainian chironomids. The high number of new records suggests that the chironomid fauna, especially from flowing waters is far from being fully discovered. Undoubtedly, it is necessary to continue the study of the chironomid fauna of the Eastern Carpathians. Particularly, the collection of the pupal exuvia could be a very useful tool in studying species richness, ecology, and distribution, but also for water quality assessment and monitoring purposes.

Acknowledgements

We gratefully acknowledge the support of the Carpathian Biosphere Reserve Administration. We are indebted to Mykola M. Voloshchuk, Peter Turis, and Milan Valachovič for providing field collection assistance. We are grateful to the reviewer and subject editor for their valuable comments to the previous version of the manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The study of mountain lakes in the Carpathians was supported by the Slovak Research and Development Agency (project No. APVV-20-0358) and Grant Agency VEGA (project No. 1/0400/21).

Author contributions

PB identified Chironomidae pupal exuviae and adults and wrote the text, MN and MS collected the data, prepared the map, tables and wrote part of the text, LH wrote part of the text.

Author ORCIDs

Milan Novikmec  <https://orcid.org/0000-0002-5192-4575>

Marek Svitok  <https://orcid.org/0000-0003-2710-8102>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Aagaard K, Solem JO, Nøst T, Hanssen O (1997) The macrobenthos of the pristine stream, Skiftesåa, Høylandet, Norway. *Hydrobiologia* 348(1/2/3): 81–94. <https://doi.org/10.1023/A:1003037200703>
- Abbou F, Fahde A (2017) Structure et diversité taxonomique des peuplements de macroinvertébrés benthiques du réseau hydrographique du bassin du Sebou (Maroc). *International Journal of Biological and Chemical Sciences* 11(4): 1785–1806. <https://doi.org/10.4314/ijbcs.v11i4.29>
- Afanasyev S, Lyashenko A, Iarochévitch A, Lietytska O, Zorina-Sakharova K, Marushevska O (2020) Pressures and impacts on ecological status of surface water bodies in Ukrainian part of the Danube River basin. In: Bănăduc D, Curtean-Bănăduc A, Pedrotti F, Cianfaglion K, Akeroyd JR (Eds) *Human Impact on Danube Watershed Biodiversity in the XXI Century. Geobotany Studies*. Springer Nature, Switzerland, 327–358. https://doi.org/10.1007/978-3-030-37242-2_16
- Ashe P, O'Connor JP (2009) A World Catalogue of Chironomidae (Diptera). Part 1. Buchonomyiinae, Chilenomyiinae, Podonominae, Aphroteniinae, Tanypodinae, Usambaromyiinae, Diamesinae, Prodiamesinae and Telfordinae. Irish Biogeographical Society and National Museum of Ireland, Dublin, 445 pp.
- Ashe P, O'Connor JP (2012) A World Catalogue of Chironomidae (Diptera). Part 2. Orthoclaadiinae. Irish Biogeographical Society and National Museum of Ireland, Dublin, 968 pp.
- Azzouzi A, Laville H, Reiss F (1992) Nouvelles récoltes de Chironomidés (Diptera) du Maroc. *Annales de Limnologie* 28(3): 225–232. <https://doi.org/10.1051/limn/1992019>
- Baranov VA (2011a) A preliminary annotated checklist of non-biting midges (Diptera, Chironomidae) of Ukraine. *Ukrainska Entomofaunistyka* 2: 7–24. <https://doi.org/10.5281/zenodo.5565484>
- Baranov VA (2011b) New and rare species of Orthoclaadiinae (Diptera, Chironomidae) from the Crimea, Ukraine. *Vestnik Zoologii* 45(5): 405–410. <https://doi.org/10.2478/v10058-011-0026-1>
- Baranov VA (2013) First records of several Tanypodinae species (Diptera, Chironomidae) from Ukraine. *Vestnik Zoologii* 47(3): 59–62. <https://doi.org/10.2478/vzoo-2013-0027>
- Baranov VA (2014) Adult female of *Chaetocladius insolitus* Caspers, 1987 (Diptera: Chironomidae) with notes on species distribution. *Aquatic Insects* 36(1): 9–14. <https://doi.org/10.1080/01650424.2015.1007071>

- Baranov VA, Ferrington Jr LC (2013) Hibernial emergence of Chironomidae in Ukraine. *Chironomus* 26(26): 33–40. <https://doi.org/10.5324/cjcr.v0i26.1616>
- Baranov VA, Przhiboro AA (2014) New records of non-biting midges (Diptera: Chironomidae) from springs and streams of the Ukrainian Carpathians (Gorgany Massif). *Zoosystematica Rossica* 23(1): 150–157. <https://doi.org/10.31610/zsr/2014.23.1.150>
- Baranov V, Lin X, Hübner J, Chimeno C (2024) Uncovering the hidden diversity of non-biting midges (Diptera, Chironomidae) from central Namibia, using morphology and DNA barcodes. *African Invertebrates* 65(1): 13–36. <https://doi.org/10.3897/afrinvertebr.65.111920>
- Bitušík P (2004) Chironomids (Diptera. Chironomidae) of the mountain lakes in the Tatra Mts. (Slovakia). A review. *Dipterologica bohemoslovaca* 12: 25–53.
- Bitušík P, Illéšová D (1998) New records of Chironomidae (Diptera, Nematocera) from Slovakia. *Biologia* 53: 238–238.
- Bitušík P, Langton PH (1994) Further new records of chironomids (Diptera. Chironomidae) from Slovakia. *Biologia* 49: 235–237.
- Bitušík P, Svitok M, Dragúňová M (2006) The actual longitudinal zonation of the river Hron (Slovakia) based on chironomid assemblages (Diptera: Chironomidae). *Acta Universitatis Carolinae. Biologica* 50: 5–17.
- Bitušík P, Novíkmec M, Hamerlik L (2020) Chironomids (Insecta, Diptera, Chironomidae) from alpine lakes in the Eastern Carpathians with comments on newly-recorded species from Ukraine. *Biodiversity Data Journal* 8: e49378. <https://doi.org/10.3897/BDJ.8.e49378>
- Borkent A (1981) The distribution and habitat preferences of the Chaoboridae (Culicomorpha: Diptera) of the Holarctic region. *Canadian Journal of Zoology* 59(1): 122–133. <https://doi.org/10.1139/z81-019>
- Borkent A (2014) World catalog of extant and fossil Chaoboridae (Diptera). *Zootaxa* 3796(3): 469–493. <https://doi.org/10.11646/zootaxa.3796.3.4>
- Calle-Martínez D, Casas JJ (2006) Chironomid species, stream classification, and water-quality assessment: The case of 2 Iberian Mediterranean mountain regions. *Journal of the North American Benthological Society* 25(2): 465–476. [https://doi.org/10.1899/0887-3593\(2006\)25\[465:CSSCAW\]2.0.CO;2](https://doi.org/10.1899/0887-3593(2006)25[465:CSSCAW]2.0.CO;2)
- Caspers N (1991) The actual biocoenotic zonation of the River Rhine exemplified by the chironomid midges (Insecta, Diptera). *Internationale Vereinigung für theoretische und angewandte Limnologie. Verhandlungen - Internationale Vereinigung für Theoretische und Angewandte Limnologie* 24(3): 1829–1834. <https://doi.org/10.1080/03680770.1989.11899080>
- Chaib N, Bouhala Z, Fouzari L, Marziali L, Samraoui B, Rossaro B (2013) Environmental factors affecting the distribution of chironomid larvae of the Seybouse wadi, North-eastern Algeria. *Journal of Limnology* 72(2): 203–214. <https://doi.org/10.4081/jlimnol.2013.e16>
- Cobo F, Soriano Ó, Báez M (2002) Chironomidae. In: Hjorth-Andersen MC-T (Ed.) *Catálogo de los Díptera de España, Portugal y Andorra*. Sociedad Entomológica Aragonesa (SEA), Zaragoza, 35–44.
- Cranston PS (1995) Introduction. In: Armitage PD, Cranston PS, Pinder LCV (Eds) *The Chironomidae: Biology and ecology of non-biting midges*. Chapman & Hall, London, 7 pp. https://doi.org/10.1007/978-94-011-0715-0_1
- de Beauvesère-Storm A, Tempelman D (2009) De dansmug *Psectrocladius schlienzi* nieuw voor Nederland, met een beschrijving van de larve (Diptera: Chironomidae). *Nederlandse Faunistische Mededelingen* 30: 17–22.

- de Jong Y (2016). Fauna Europaea. Fauna Europaea Consortium. Checklist dataset. <https://doi.org/10.15468/ymk1bx> [GBIF.org on 2024-03-21]
- Didenko A, Volikov Y, Baranov V, Kruzhylina S, Gurbyk A, Bielikova O (2021) Chironomid diversity in the diets of Ponto-Caspian gobiids in a freshwater habitat: Implications for resource partitioning. *Limnologica* 89: 125890. <https://doi.org/10.1016/j.limno.2021.125890>
- Dittmar H (2012) Beiträge zur aquatischen Insektenfauna des Sauerlandes. Abhandlungen aus dem Westfälischen Museum für Naturkunde 74: 3–41.
- Eggermont H, Heiri O (2012) The chironomid-temperature relationship: Expression in nature and palaeoenvironmental implications. *Biological Reviews of the Cambridge Philosophical Society* 87(2): 430–456. <https://doi.org/10.1111/j.1469-185X.2011.00206.x>
- Ekrem T (2004) Immature stages of European *Tanytarsus* species I. The *eminulus*-, *gregarius*-, *lugens*- and *mendax* species groups (Diptera, Chironomidae). *Deutsche Entomologische Zeitschrift*. *Deutsche Entomologische Zeitschrift* 51(1): 97–146. <https://doi.org/10.1002/mmnd.20040510110>
- Epler JH, Ekrem T, Cranston PS (2013) 10. The larvae of Chironominae (Diptera: Chironomidae) of the Holarctic Region – Keys and diagnoses. In: Andersen T, Sæther OA, Cranston PS & Epler JH (Eds) *Chironomidae of the Holarctic Region. Keys and diagnoses. Larvae*. *Insect Systematics & Evolution, Suppl.* 66: 387–556.
- Ferrington Jr LC (2007) Global diversity of non-biting midges (Chironomidae; Insecta-Diptera) in freshwater. *Hydrobiologia* 595(1): 447–455. <https://doi.org/10.1007/s10750-007-9130-1>
- Gandouin E, Maasri A, Van Vliet-Lanoe B, Franquet E (2006) Chironomid (Insecta: Diptera) assemblages from a gradient of lotic and lentic waterbodies in river floodplains of France: a methodological tool for paleoecological applications. *Journal of Paleolimnology* 35(1): 149–166. <https://doi.org/10.1007/s10933-005-8149-4>
- Giłka W (2007) A faunistic review of chironomids of the tribe Tanytarsini (Diptera: Chironomidae) of the Tatra National Park. *Dipteron* 23: 11–17.
- Golubets MA (1977) Eastern-Carpathian Mountain subprovince. In: Barbarych AI (Ed.) *Geobotanical regionalization of Ukrainian SSR*. Naukova dumka, Kyiv, 18–44.
- Hayford B (2005) New records of Chironomidae (Insecta: Diptera) from Mongolia with review of distribution and biogeography of Mongolian Chironomidae. *Journal of the Kansas Entomological Society* 78(2): 192–200. <https://doi.org/10.2317/0406.24.1>
- Hirvenoja M, Moubayed J (1989) The adults and preimaginal stages of continental populations of *Cricotopus* (*Cricotopus*) *beckeri* Hirvenoja (Diptera, Chironomidae). *Annales de Limnologie* 25(3): 251–254. <https://doi.org/10.1051/limn/1989027>
- Ivanik O, Shevchuk V, Kravchenko D, Shpyrko S, Yanchenko V, Gadiatska K (2019) Geological and geomorphological factors of natural hazards in Ukrainian Carpathians. *Journal of Ecological Engineering* 20(4): 177–186. <https://doi.org/10.12911/22998993/102964>
- Joniak T, Domek P (2006) Influence of humification on biodiversity of lake benthic macroinvertebrates. *Acta Agrophysica* 7: 3636–368.
- Kettani K, Langton PH (2012) Les Chironomidae du Maroc (Diptera, Nematocera). *Bulletin de la Société Entomologique de France* 117(4): 411–424. <https://doi.org/10.3406/bsef.2012.3065>
- Klink A (2010) Macroinvertebrates of the Seine basin. *Hydrobiologisch Adviesburo Klink Rapporten en Mededelingen* 108, Wageningen, 71 pp.
- Kobiv Y (2017) Response of rare alpine plant species to climate change in the Ukrainian Carpathians. *Folia Geobotanica* 52(2): 217–226. <https://doi.org/10.1007/s12224-016-9270-z>

- Kownacki A (1985) Spring benthic macroinvertebrate communities of selected streams in the High Caucasus (Azerbaijan SSR). *Hydrobiologia* 123(2): 137–144. <https://doi.org/10.1007/BF00018975>
- Krashennnikov AB (2012) Phenology of some chironomid species (Diptera, Chironomidae) of the Middle Urals. *Fauna Norvegica* 31: 55–55. <https://doi.org/10.5324/fn.v31i0.1371>
- Kresonja M (2018) Urban aquatic habitats, neglected sources of biodiversity. Comparison of chironomid communities (Chironomidae, Diptera) in fountains, artificial lakes and the River Drava. PhD Thesis, University Josip Juraj Strossmayer in Osijek, Osijek, Croatia.
- Kuper JT, Verberk WCEP (2011) Fenologie, habitat en verspreiding van Pluimmuggen in Nederlandse hoogvenen (Diptera: Chaoboridae). *Nederlandse Faunistische Mededelingen* 35: 73–82.
- Langton PH (1991) A key to pupal exuviae of West Palaearctic Chironomidae. Privately published. Huntington, Cambridgeshire, England, 368 pp.
- Langton PH, Orendt C (1996) A collection of Chironomidae from La Gomera, Canary Islands. *Entomologist's Gazette* 47: 279–281.
- Langton PH, Pinder LCV (2007a) Keys to the adult male Chironomidae of Britain and Ireland 1. Scientific Publication - Freshwater Biological Association 64: 1–239.
- Langton PH, Pinder LCV (2007b) Keys to the adult male Chironomidae of Britain and Ireland 2. Scientific Publication - Freshwater Biological Association 64: 1–168.
- Langton PH, Visser H (2003) Chironomidae exuvia. A key to pupal exuvia of the West Palaearctic Region. Interactive Identification System for the European Limnofauna (IISEL), World Biodiversity Database, CD-ROM Series.
- Langton PH, Bitusik P, Mitterova J (2013) A contribution towards a revision of West Palaearctic *Procladius* Skuse (Diptera: Chironomidae). *Chironomus* 26(26): 41–44. <https://doi.org/10.5324/cjcr.v0i26.1620>
- Laville H, Langton P (2002) The lotic Chironomidae (Diptera) of Corsica (France). *Annales de Limnologie* 38(1): 53–64. <https://doi.org/10.1051/limn/2002006>
- Laville H, Reiss F (1992) The chironomid fauna of the Mediterranean region reviewed. *Netherlands Journal of Aquatic Ecology* 26(2–4): 239–245. <https://doi.org/10.1007/BF02255247>
- Laville H, Tourenq JN (1968) Nouvelles récoltes de chironomides en Camargue et dans les Marismas du Guadalquivir (Diptères). *Annales de Limnologie* 4(1): 73–80. <https://doi.org/10.1051/limn/1968010>
- Laville H, Vinçon G (1986) Inventaire 1986 des Chironomidés (Diptera) connus des Pyrénées. *Annales de Limnologie* 22(3): 239–251. <https://doi.org/10.1051/limn/1986022>
- Laville H, Vinçon G (1991) A typological study of Pyrenean streams: Comparative analysis of the Chironomidae (Diptera) communities in the Ossau and Aure Valleys. *Internationale Vereinigung für theoretische und angewandte Limnologie. Verhandlungen - Internationale Vereinigung für Theoretische und Angewandte Limnologie* 24(3): 1775–1784. <https://doi.org/10.1080/03680770.1989.11899070>
- Lencioni V, Marziali L, Rossaro B (2011) Diversity and distribution of chironomids (Diptera, Chironomidae) in pristine Alpine and pre-Alpine springs (Northern Italy). *Journal of Limnology* 70(1s): 106–121. <https://doi.org/10.4081/jlimnol.2011.s1.106>
- Luoto T, Nevalainen L (2009) Larval chaoborid mandibles in surface sediments of small shallow lakes in Finland: Implications for palaeolimnology. *Hydrobiologia* 631(1): 185–195. <https://doi.org/10.1007/s10750-009-9810-0>

- MacKay NA, Carpenter SR, Soranno PA, Vanni MJ (1990) The impact of two *Chaoborus* species on a zooplankton community. *Canadian Journal of Zoology* 68(5): 981–985. <https://doi.org/10.1139/z90-141>
- Martínez DC, Quero AV, Jiménez JC (1995) Les chironomidés (Diptera) du bassin du Haut-Guadalquivir (Sierra de Cazorla, sud de l'Espagne). *Annales de Limnologie* 31(3): 201–213. <https://doi.org/10.1051/limn/1995018>
- Moller Pillot HKM (2009) Chironomidae larvae of the Netherlands and adjacent low-lands: biology and ecology of the Chironomini. KNNV Publishing, Zeist, 270 pp.
- Moller Pillot HKM (2013) Chironomidae Larvae of the Netherlands and Adjacent Low-lands. Biology and Ecology of the aquatic Orhocladiinae. KNNV Publishing, Zeist, 312 pp.
- Molnár A, Végvári Z, Tóthmérész B (2017) Identification of floral relicts based on spatial distance of isolation. *Forests* 8(11): 459. <https://doi.org/10.3390/f8110459>
- Móra A, Tóth M, Debreceni Á, Csépes E (2006) Contribution to the chironomid fauna (Diptera: Chironomidae) of the Upper-Tisza, NE Hungary. *Folia Historico-naturalia Musei Matraensis* 30: 253–261.
- Móra A, Polyák L, Farkas A (2013) Contribution to the Chironomidae (Diptera) fauna of the Sajó/Slaná River, Hungary and Slovakia. *Acta Biologica Debrecina-Supplementum Oecologica Hungarica* 31: 69–81.
- Moubayed J, Bernard A, Claude J, Decoin R, Tissot B (2019) Inventaire des Chironomidae de la réserve naturelle nationale du Lac de Remoray. II. Liste des espèces recensées en 2019 avec commentaires sur leur écologie et leur distribution géographique (Diptera). *Ephemera* 20: 113–131.
- Moubayed-Breil J (2008) Non-biting midges from Continental France: New records, faunal and biogeographical outline (Diptera, Chironomidae). *Ephemera* 9: 17–32.
- Moubayed-Breil J, Ashe P (2016) New records and additions to the database on the geographical distribution of some threatened chironomid species from continental France (Diptera, Chironomidae). *Ephemera* 16: 93–108.
- Moubayed-Breil J, Baranov VA (2018) Taxonomic notes on the genus *Hydrobaenus* with description of *H. simferopolus* sp. nov. from Crimea (Diptera: Chironomidae). *Acta Entomologica Musei Nationalis Pragae* 58(2): 347–355. <https://doi.org/10.2478/aemnp-2018-0029>
- Moubayed-Breil J, Langton PH, Ashe P (2012) *Rheotanytarsus dactylophoreus*, a new mountain species from streams in the Eastern Pyrenees and Corsica (Diptera: Chironomidae). *Fauna Norvegica* 31: 167–167. <https://doi.org/10.5324/fn.v31i0.1375>
- Nicacio G, Juen L (2015) Chironomids as indicators in freshwater ecosystems: An assessment of the literature. *Insect Conservation and Diversity* 8(5): 393–403. <https://doi.org/10.1111/icad.12123>
- Nilssen JP (1974) On the ecology and distribution of the Norwegian larvae of *Chaoborus* (Diptera, Chaoboridae). *Norsk Entomologisk Tidsskrift* 1: 37–44.
- Novikoff A, Hurdu BI (2015) A critical list of endemic vascular plants in the Ukrainian Carpathians. *Contributii Botanice* 50: 43–91.
- Orendt C (2002a) Die Chironomidenfauna des Inns bei Mühldorf (Oberbayern). *Lauterbornia* 44: 109–120.
- Orendt C (2002b) Biozönotische Klassifizierung naturnaher Flussabschnitte des nördlichen Alpenvorlandes auf der Grundlage der Zuckmücken-Lebensgemeinschaften (Diptera: Chironomidae). *Lauterbornia* 44: 121–146.
- Orendt C, Reiff N (2003) Rote Liste gefährdeter Zuckmücken (Diptera: Chironomidae) Bayerns. *Bayerisches Landesamt für Umweltschutz* 166: 301–304.

- Paasivirta L (2014) Checklist of the family Chironomidae (Diptera) of Finland. ZooKeys 441: 63–90. <https://doi.org/10.3897/zookeys.441.7461>
- Pape T, Blagoderov V, Mostovski MB (2011) Order Diptera Linnaeus, 1758. In: Zhang ZO (Ed.) Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 222–229. <https://doi.org/10.11646/zootaxa.3148.1.42>
- Parma S (1969) Notes on the larval taxonomy, ecology, and distribution of the Dutch *Chaoborus* species (Diptera, Chaoboridae). Beaufortia 17: 21–50.
- Pozdeev IV (2012) A new data on Chironomid fauna (Diptera, Chironomidae) of the Upper and Middle Kama River basin. Evraziatskii Entomologicheskii Zhurnal 11: 87–90.
- Prat N, Puntí T, Rieradevall M (2016) The use of larvae and pupal exuviae to study the biodiversity of Chironomidae in Mediterranean streams. Journal of Entomological and Acarological Research 48 (1): 29–36. <https://doi.org/10.4081/jear.2016.5529>
- Reiss F (1986) A contribution to the zoogeography of the Turkish Chironomidae (Diptera). Israel Journal of Entomology 19: 161–170.
- Resh V, Rosenberg D (2008) Water pollution and insects. In: Capinera J (Ed.) Encyclopedia of Entomology, Springer Science+Business Media B.V., Dordrecht, 4158–4168.
- Ringe F (1974) Chironomiden-Emergenz 1970 in Breitenbach und Rohrwiesenbach. Schlitzer Produktionsbiologische Studies (10). Archiv für Hydrobiologie (Supplementband 45): 212–304.
- Rossaro B (1991) Chironomids and water temperature. Aquatic Insects 13(2): 87–98. <https://doi.org/10.1080/01650429109361428>
- Rossaro B, Lencioni V (2015) A key to larvae of *Diamesa* Meigen, 1835 (Diptera, Chironomidae), well known as adult males and pupae from Alps (Europe). Journal of Entomological and Acarological Research 47(3): 123–138. <https://doi.org/10.4081/jear.2015.5516>
- Rossaro B, Lencioni V, Casalegno C (2003) Revision of West Palaearctic species of *Orthocladius* s. str. van der Wulp, 1874 (Diptera: Chironomidae: Orthoclaadiinae), with a new key to species. Studi Trentini di Scienze Naturali. Acta Biologica 79: 213–241.
- Ruse LP (1995) Chironomid community structure deduced from larvae and pupal exuvia of a chalk stream. Hydrobiologia 315(2): 135–142. <https://doi.org/10.1007/BF00033625>
- Sæther OA (1997) Family Chaoboridae. In: Papp L, Darvas B (Eds) Contributions to a Manual of Palaearctic Diptera 2. Nematocera and Lower Brachycera. Science Herald, Budapest, 305–323.
- Sæther OA, Andersen T (2013) 7. The larvae of Diamesinae (Diptera: Chironomidae) of the Holarctic Region – Keys and diagnoses. In: Andersen T, Sæther OA, Cranston PS, Epler JH (Eds) Chironomidae of the Holarctic Region. Keys and diagnoses. Larvae. Insect Systematics & Evolution, Suppl. 66: 145–178.
- Sæther OA, Ashe P, Murray DA (2000) Family Chironomidae. In: Papp L, Darvas B (Eds) Contribution to a Manual of Palaearctic Diptera (with special reference to flies of economic importance). Appendix. Science Herald, Budapest, 113–334.
- Schöll F, Haybach A (2004) Typology of large European rivers according to their Chironomidae communities (Insecta: Diptera). Annales de Limnologie-International Journal of Limnology 40: 309–316). <https://doi.org/https://doi.org/10.1051/limn/2004028>
- Sealock AW, Ferrington LC (2008) Key and descriptions of the Chironomidae pupal exuvia of Hardwood Creek, near Hugo, Minnesota. Chironomidae Research Group, University of Minnesota, St. Paul, 86 pp.
- Sołtys-Lelek A, Barabasz-Krasny B, Możdżeń K (2014) Selected aspects of the protection of biotopes on the example of the Ojców National Park (Southern Poland). Annales Universitatis Paedagogicae Cracoviensis 174: 98–117.

- Soriano O, Cobo F (2006) Lista faunística de los quironómidos (Diptera, Chironomidae) de Madrid (España). *Graellsia* 62(Extra): 7–20. <https://doi.org/10.3989/graelisia.2006.v62.iExtra.107>
- Spies M, Sæther OA (2013) Fauna Europaea: Chironomidae. In: Pape T, Beuk P (Eds) Fauna Europaea: Diptera. Fauna Europaea version 2017.06. <https://www.faunaeur.org> [accessed 12 September 2019]
- Stenson JA (1981) The role of predation in the evolution of morphology, behaviour and life history of two species of *Chaoborus*. *Oikos* 37(3): 323–327. <https://doi.org/10.2307/3544123>
- Stur E, Ekrem T (2006) A revision of West Palaearctic species of the *Micropsectra atrofasciata* group (Diptera: Chironomidae). *Zoological Journal of the Linnean Society* 146(2): 165–225. <https://doi.org/10.1111/j.1096-3642.2006.00198.x>
- Syrovátka V, Langton PH (2015) First records of *Lasiodiamesa gracilis* (Kieffer, 1924), *Parochlus kiefferi* (Garrett, 1925) and several other Chironomidae from the Czech Republic and Slovakia. *Chironomus* 28(28): 45–56. <https://doi.org/10.5324/cjcr.v0i28.1953>
- Tatole V (2023) An updated checklist of chironomid species (Diptera, Chironomidae) from Romania. *Travaux du Museum National d'Histoire Naturelle "Grigore Antipa" / Muzeul National de Istorie Naturala "Grigore Antipa"* 66(1): 17–105. <https://doi.org/10.3897/travaux.66.e102065>
- Tolonen KT, Brodersen KP, Kleisborg TA, Holmgren K, Dahlberg M, Hamerlik L, Hämäläinen H (2012) Phantom midge-based models for inferring past fish abundances. *Journal of Paleolimnology* 47(4): 531–547. <https://doi.org/10.1007/s10933-012-9579-4>
- Tourenq JN (1976) Recherches écologiques sur les chironomides (Diptera) de Camargue. I. Etude faunistique et biogéographique. *Annales de Limnologie* 12(1): 17–74. <https://doi.org/10.1051/limn/1976009>
- Vallenduuk HJ, Moller Pillot HKM (2007) Chironomidae larvae of the Netherlands and adjacent lowlands. General ecology and Tanypodinae. KNNV Publishing, Zeist, 144 pp.
- Vyshnevskyi VI, Donich OA (2021) Climate change in the Ukrainian Carpathians and its possible impact on river runoff. *Acta Hydrologica Slovaca* 22(1): 3–14. <https://doi.org/10.31577/ahs-2021-0022.01.0001>

Supplementary material 1

Checklist of chironomid and chaoborid taxa recorded in studied streams and lakes of the Ukrainian part of the Eastern Carpathians

Authors: Peter Bitušík, Milan Novikmec, Marek Svitok, Ladislav Hamerlík

Data type: xlsx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1211.125436.suppl1>